



Please cancel claims 66 and 67 without prejudice.

Kindly amend the following claims 1, 19, 26, 32, 52, 54, 57, 64 and 68 as follows.

 1. (Amended) A peptide conjugate comprising X and Z,

wherein X is a pharmacologically active heteropolymeric peptide sequence, and

 wherein Z is a stabilising peptide sequence, of 4-20 amino acid units covalently bonded by its N terminus to the C terminus end of X wherein each amino acid unit in said stabilising peptide sequence, Z is selected from the group consisting of Ala, Leu, Ser, Thr, Tyr, Asn, Gln, Asp, Glu, Lys, Arg, His, Met, Orn, and amino acid units of the general formula I



wherein R¹ and R² are selected from the group consisting of hydrogen, C₁₋₆-alkyl, phenyl, and phenyl-methyl, wherein C₁₋₆-alkyl is optionally substituted with from one to three substituents selected from halogen, hydroxy, amino, cyano, nitro, sulfono, and carboxy, and phenyl and phenyl-methyl is optionally substituted with from one to three substituents selected from C₁₋₆-alkyl, C₂₋₆-alkenyl, halogen, hydroxy, amino, cyano, nitro, sulfono, and carboxy, or R¹ and R² together with the carbon atom to which they are bound form a cyclopentyl, cyclohexyl, or cycloheptyl ring; and


wherein the ratio between the half-life of said peptide conjugate and the half-life of the corresponding pharmacologically active peptide sequence, X, when treated with carboxypeptidase A or leucine aminopeptidase in about 50 mM phosphate buffer solution at

ES conclude
about pH 7.4 at about 37°C or in serum or plasma is at least 2; or a salt thereof,
wherein Z comprises at least two identical amino acid units.

19. (Amended) A peptide conjugate according to claim 1,
wherein Z is Lys-Lys-Lys-Lys (SEQ ID NO. 55), Xaa-Lys-Lys-Lys,
Lys-Xaa-Lys-Lys, Lys-Lys-Xaa-Lys, Lys-Lys-Lys-Xaa, Xaa-Xaa-Lys-
Lys, Xaa-Lys-Xaa-Lys, Xaa-Lys-Lys-Xaa, Lys-Xaa-Xaa-Lys, Lys-Xaa-
Lys-Xaa, Lys-Lys-Xaa-Xaa, Xaa-Xaa-Xaa-Lys, Xaa-Xaa-Lys-Xaa, Xaa-
Lys-Xaa-Xaa, Lys-Xaa-Xaa-Xaa, Xaa-Xaa-Xaa-Xaa, Lys-Lys-Lys-Lys-
Lys (SEQ ID NO. 56), Xaa-Lys-Lys-Lys-Lys (SEQ ID NO. 57), Lys-
Xaa-Lys-Lys-Lys (SEQ ID NO. 58), Lys-Lys-Xaa-Lys-Lys (SEQ ID NO.
59), Lys-Lys-Lys-Xaa-Lys (SEQ ID NO. 60), Lys-Lys-Lys-Lys-Xaa
(SEQ ID NO. 61), Xaa-Xaa-Lys-Lys-Lys, Xaa-Lys-Xaa-Lys-Lys, Xaa-
Lys-Lys-Xaa-Lys, Xaa-Lys-Lys-Lys-Xaa, Lys-Xaa-Xaa-Lys-Lys, Lys-
Xaa-Lys-Xaa-Lys, Lys-Xaa-Lys-Lys-Xaa, Lys-Lys-Xaa-Xaa-Lys, Lys-
Lys-Xaa-Lys-Xaa, Lys-Lys-Lys-Xaa-Xaa, Lys-Lys-Xaa-Xaa-Xaa, Lys-
Xaa-Lys-Xaa-Xaa, Lys-Xaa-Xaa-Lys-Xaa, Lys-Xaa-Xaa-Xaa-Lys, Xaa-
Lys-Lys-Xaa-Xaa, Xaa-Lys-Xaa-Xaa-Lys, Xaa-Xaa-Lys-Lys-Xaa, Xaa-
Xaa-Lys-Xaa-Lys, Xaa-Xaa-Xaa-Lys-Lys, Lys-Xaa-Xaa-Xaa-Xaa, Xaa-
Lys-Xaa-Xaa-Xaa, Xaa-Xaa-Lys-Xaa-Xaa, Xaa-Xaa-Xaa-Lys-Xaa, Xaa-
Xaa-Xaa-Xaa-Lys, Xaa-Xaa-Xaa-Xaa-Xaa, Lys-Lys-Lys-Lys-Lys-Lys
(SEQ ID NO. 62), Xaa-Lys-Lys-Lys-Lys-Lys (SEQ ID NO. 63), Lys-
Xaa-Lys-Lys-Lys-Lys (SEQ ID NO. 64), Lys-Lys-Xaa-Lys-Lys-Lys
(SEQ ID NO. 65), Lys-Lys-Lys-Xaa-Lys-Lys (SEQ ID NO. 66), Lys-
Lys-Lys-Lys-Xaa-Lys (SEQ ID NO. 67), Lys-Lys-Lys-Lys-Lys-Xaa
(SEQ ID NO. 68), Xaa-Xaa-Lys-Lys-Lys-Lys (SEQ ID NO. 69), Xaa-
Lys-Xaa-Lys-Lys-Lys (SEQ ID NO. 70), Xaa-Lys-Lys-Xaa-Lys-Lys
(SEQ ID NO. 71), Xaa-Lys-Lys-Lys-Xaa-Lys (SEQ ID NO. 72), Xaa-
Lys-Lys-Lys-Lys-Xaa (SEQ ID NO. 73), Lys-Xaa-Xaa-Lys-Lys-Lys
(SEQ ID NO. 74), Lys-Xaa-Lys-Xaa-Lys-Lys (SEQ ID NO. 75), Lys-

12
Conf.

Xaa-Lys-Lys-Xaa-Lys (SEQ ID NO. 76), Lys-Xaa-Lys-Lys-Lys-Xaa
(SEQ ID NO. 77), Lys-Lys-Xaa-Xaa-Lys-Lys (SEQ ID NO. 78), Lys-
Lys-Xaa-Lys-Xaa-Lys (SEQ ID NO. 79), Lys-Lys-Xaa-Lys-Lys-Xaa
(SEQ ID NO. 80), Lys-Lys-Lys-Xaa-Xaa-Lys (SEQ ID NO. 81), Lys-
Lys-Lys-Xaa-Lys-Xaa (SEQ ID NO. 82), Lys-Lys-Lys-Lys-Xaa-Xaa
(SEQ ID NO. 83), Xaa-Xaa-Xaa-Lys-Lys-Lys, Xaa-Xaa-Lys-Xaa-Lys-
Lys, Xaa-Xaa-Lys-Lys-Xaa-Lys, Xaa-Xaa-Lys-Lys-Lys-Xaa, Xaa-Lys-
Xaa-Xaa-Lys-Lys, Xaa-Lys-Xaa-Lys-Xaa-Lys, Xaa-Lys-Xaa-Lys-Lys-
Xaa, Xaa-Lys-Lys-Xaa-Xaa-Lys, Xaa-Lys-Lys-Xaa-Lys-Xaa, Xaa-Lys-
Lys-Lys-Xaa-Xaa, Lys-Lys-Lys-Xaa-Xaa-Xaa, Lys-Lys-Xaa-Lys-Xaa-
Xaa, Lys-Lys-Xaa-Xaa-Lys-Xaa, Lys-Lys-Xaa-Xaa-Xaa-Lys, Lys-Xaa-
Lys-Lys-Xaa-Xaa, Lys-Xaa-Lys-Xaa-Lys-Xaa, Lys-Xaa-Lys-Xaa-Xaa-
Lys, Lys-Xaa-Xaa-Lys-Lys-Xaa, Lys-Xaa-Xaa-Lys-Xaa-Lys, Lys-Xaa-
Xaa-Xaa-Lys-Lys, Lys-Lys-Xaa-Xaa-Xaa-Xaa, Lys-Xaa-Lys-Xaa-Xaa-
Xaa, Lys-Xaa-Xaa-Lys-Xaa-Xaa, Lys-Xaa-Xaa-Xaa-Lys-Xaa, Lys-Xaa-
Xaa-Xaa-Xaa-Lys, Xaa-Lys-Lys-Xaa-Xaa-Xaa, Xaa-Lys-Xaa-Lys-Xaa-
Xaa, Xaa-Lys-Xaa-Xaa-Lys-Xaa, Xaa-Lys-Xaa-Xaa-Xaa-Lys, Xaa-Xaa-
Lys-Lys-Xaa-Xaa, Xaa-Xaa-Lys-Xaa-Lys-Xaa, Xaa-Xaa-Lys-Xaa-Xaa-
Lys, Xaa-Xaa-Xaa-Lys-Lys-Xaa, Xaa-Xaa-Xaa-Lys-Lys-Lys, Xaa-Xaa-
Xaa-Lys-Xaa-Lys, Xaa-Xaa-Xaa-Xaa-Lys-Lys, Lys-Xaa-Xaa-Xaa-Xaa-
Xaa, Xaa-Lys-Xaa-Xaa-Xaa-Xaa, Xaa-Xaa-Lys-Xaa-Xaa-Xaa, Xaa-Xaa-
Xaa-Lys-Xaa-Xaa, Xaa-Xaa-Xaa-Xaa-Lys-Xaa, Xaa-Xaa-Xaa-Xaa-Xaa-
Lys, or Xaa-Xaa-Xaa-Xaa-Xaa-Xaa, wherein each Xaa is
independently selected from the group consisting of Ala, Leu,
Ser, Thr, Tyr, Asn, Gln, Asp, Glu, Arg, His, Met, Orn, and amino
acids of the formula I



(I)


E2
conclude

wherein R^1 and R^2 are selected from the group consisting of hydrogen, C_{1-6} -alkyl, phenyl, and phenyl-methyl, wherein C_{1-6} -alkyl is optionally substituted with from one to three substituents selected from halogen, hydroxy, amino, cyano, nitro, sulfono, and carboxy, and phenyl and phenyl-methyl is optionally substituted with from one to three substituents selected from C_{1-6} -alkyl, C_{2-6} -alkenyl, halogen, hydroxy, amino, cyano, nitro, sulfono, and carboxy, or R^1 and R^2 together with the carbon atom to which they are bonded form a cyclopentyl, cyclohexyl, or cycloheptyl ring.

E3


26. (Amended) A method for the preparation of a peptide conjugate (X-Z) as defined in claim 2, comprising the steps of:

- a) coupling an N- α -protected amino acid or N- α -protected dipeptide in the carboxyl activated form, in the C-terminal activated form to an immobilised peptide sequence H-Z-SSM, thereby forming an immobilised N- α -protected peptide fragment,
 - b) removing the N- α -protecting group, thereby forming an immobilised peptide fragment having an unprotected N-terminal end,
 - c) coupling an additional N- α -protected amino acid in the carboxyl activated form, or an additional N- α -protected dipeptide in the C-terminal activated form to the N-terminal end of the immobilised peptide fragment, and repeating the removal/coupling step procedure in step b) and
 - c) until the desired peptide sequence X is obtained, and then
 - d) cleaving off the peptide conjugate from the solid support material.
-

 32. (Amended) A composition comprising a peptide conjugate according to claim 1, and a pharmaceutical acceptable carrier.

 52. (Amended) A peptide conjugate comprising X and Z,

wherein X is a pharmacologically active heteropolymeric peptide sequence, and

 wherein Z is a stabilising peptide sequence of 4-20 amino acid units covalently bound by its N terminus to the C terminus end of X, wherein each amino acid unit in said stabilising peptide sequence Z is selected from the group consisting of Ala, Leu, Ser, Thr, Tyr, Asn, Gln, Asp, Glu, Lys, Arg, His, Met, Orn, and amino acid units of the general formula I



wherein R¹ and R² are selected from the group consisting of hydrogen, C₁₋₆-alkyl, phenyl, and phenyl-methyl, wherein C₁₋₆-alkyl is optionally substituted with from one to three substituents selected from halogen, hydroxy, amino, cyano, nitro, sulfono, and carboxy, and phenyl and phenyl-methyl is optionally substituted with from one to three substituents selected from C₁₋₆-alkyl, C₂₋₆-alkenyl, halogen, hydroxy, amino, cyano, nitro, sulfono, and carboxy, or R¹ and R² together with the carbon atom to which they are bound form a cyclopentyl, cyclohexyl, or cycloheptyl ring; and

wherein the ratio between the half-life of said peptide conjugate and the half-life of the corresponding pharmacologically active peptide sequence X, when treated with carboxypeptidase A or leucine aminopeptidase in about 50 mM phosphate buffer solution at about pH 7.4 at about 37°C or in serum or plasma is at least 3; or a salt thereof, wherein Z comprises at least two identical amino acid units.

54. (Amended) A peptide conjugate comprising X and Z,

wherein X is a pharmacologically active heteropolymeric peptide sequence, and

wherein Z is a stabilising peptide sequence of 4-10 amino acid units covalently bound by its N terminus to the C terminus end of X, wherein each amino acid unit in said stabilising peptide sequence Z is selected from the group consisting of Ala, Leu, Ser, Thr, Tyr, Asn, Gln, Asp, Glu, Lys, Arg, His, Met, Orn, and amino acid units of the general formula I




wherein R^1 and R^2 are selected from the group consisting of hydrogen, C_{1-6} -alkyl, phenyl, and phenyl-methyl, wherein C_{1-6} -alkyl is optionally substituted with from one to three substituents selected from halogen, hydroxy, amino, cyano, nitro, sulfono, and carboxy, and phenyl and phenyl-methyl is optionally substituted with from one to three substituents selected from C_{1-6} -alkyl, C_{2-6} -alkenyl, halogen, hydroxy, amino, cyano, nitro, sulfono, and carboxy, or R^1 and R^2 together with the carbon atom to which they are bound form a cyclopentyl, cyclohexyl, or cycloheptyl ring; and

wherein the ratio between the half-life of said peptide conjugate and the half-life of the corresponding pharmacologically active peptide sequence X, when treated with carboxypeptidase A or leucine aminopeptidase in about 50 mM phosphate buffer solution at about pH 7.4 at about 37°C or in serum or plasma is at least 2; or a salt thereof, wherein Z comprises at least two identical amino acid units.

57. (Amended) A peptide conjugate comprising X and Z,

wherein X is a pharmacologically active heteropolymeric peptide sequence, and

wherein Z is a stabilising peptide sequence of 4-20 amino acid units covalently bound by its N terminus to the C terminus end of X, wherein each amino acid unit in said stabilising peptide sequence Z is selected from the group consisting of Ala, Leu, Ser, Thr, Tyr, Asn, Gln, Asp, Glu, Lys, Arg, His, Met, Orn, and amino acid units of the general formula I



wherein R¹ and R² are selected from the group consisting of hydrogen, C₁₋₆-alkyl, phenyl, and phenyl-methyl, wherein C₁₋₆-alkyl is optionally substituted with from one to three substituents selected from halogen, hydroxy, amino, cyano, nitro, sulfono, and carboxy, and phenyl and phenyl-methyl is optionally substituted with from one to three substituents selected from C₁₋₆-alkyl, C₂₋₆-alkenyl, halogen, hydroxy, amino, cyano, nitro, sulfono, and carboxy, or R¹ and R² together with the carbon atom to which they are bound form a cyclopentyl, cyclohexyl, or cycloheptyl ring; and

wherein the ratio between the half-life of said peptide conjugate and the half-life of the corresponding pharmacologically active peptide sequence X, when treated with carboxypeptidase A or leucine aminopeptidase in about 50 mM phosphate buffer solution at about pH 7.4 at about 37°C or in serum or plasma is at least [about] 2; or a salt thereof, wherein Z comprises at least two or three Lys amino acid units.

64. (Amended) A peptide conjugate comprising X and Z,

wherein X is a pharmacologically active heteropolymeric peptide sequence, and

wherein Z is a stabilising peptide sequence of 4-20 amino acid units covalently bound by its N terminus to the C terminus end of X, wherein each amino acid unit in said stabilising peptide sequence Z is selected from the group consisting of Ala, Leu, Ser, Thr, Tyr, Asn, Gln, Asp, Glu, Lys, Arg, His, Met, Orn, and amino acid units of the general formula I



wherein R¹ and R² are selected from the group consisting of hydrogen, C₁₋₆-alkyl, phenyl, and phenyl-methyl, wherein C₁₋₆-alkyl is optionally substituted with from one to three substituents selected from halogen, hydroxy, amino, cyano, nitro, sulfono, and carboxy, and phenyl and phenyl-methyl is optionally substituted with from one to three substituents selected from C₁₋₆-alkyl, C₂₋₆-alkenyl, halogen, hydroxy, amino, cyano, nitro, sulfono, and carboxy, or R¹ and R² together with the carbon atom to which they are bound form a cyclopentyl, cyclohexyl, or cycloheptyl ring; and

wherein the ratio between the half-life of said peptide conjugate and the half-life of the corresponding pharmacologically active peptide sequence X, when treated with carboxypeptidase A or leucine aminopeptidase in about 50 mM phosphate buffer solution at about pH 7.4 at about 37°C or in serum or plasma is at least 2; or a salt thereof, wherein said pharmacologically active peptide sequence (X) consists of at the most about 65 amino acid units, wherein Z comprises at least two identical amino acid units.

EG
2/2/01

68. (Amended) A peptide conjugate comprising X and Z,

wherein X is a pharmacologically active heteropolymeric peptide sequence, and

wherein Z is a stabilising peptide sequence of 4-20 amino acid units covalently bound by its N terminus to the C terminus end of X, wherein each amino acid unit in said stabilising peptide sequence Z is selected from the group consisting of Ala, Leu, Ser, Thr, Tyr, Asn, Gln, Asp, Glu, Lys, Arg, His, Met, Orn, and amino acid units of the general formula I



wherein R^1 and R^2 are selected from the group consisting of hydrogen, C_{1-6} -alkyl, phenyl, and phenyl-methyl, wherein C_{1-6} -alkyl is optionally substituted with from one to three substituents selected from halogen, hydroxy, amino, cyano, nitro, sulfono, and carboxy, and phenyl and phenyl-methyl is optionally substituted with from one to three substituents selected from C_{1-6} -alkyl, C_{2-6} -alkenyl, halogen, hydroxy, amino, cyano, nitro, sulfono, and carboxy, or R^1 and R^2 together with the carbon atom to which they are bound form a cyclopentyl, cyclohexyl, or cycloheptyl ring; and

wherein the ratio between the half-life of said peptide conjugate and the half-life of the corresponding pharmacologically active peptide sequence X, when treated with carboxypeptidase A or leucine aminopeptidase in about 50 mM phosphate buffer solution at about pH 7.4 at about 37°C or in serum or plasma is at least [about] 2; or a salt thereof,

wherein,

Z is $\text{Lys}_p\text{-Xaa}_q$ or $\text{Xaa}_p\text{-Lys}_q$, wherein p and q are integers in the range from 1 to 14, with the proviso that $p+q$ is in the range of 3-15, and each Xaa is independently selected from the group consisting of Ser, Thr, Tyr, Asn, Gln, Asp, Glu, Arg, His, Orn, 2,4-diaminobutanoic acid, 2,3-diaminopropanoic acid and Met,

and further wherein,

X is selected from the group consisting of AF 12505 (Ile-Glu-Gly-Pro-Thr-Leu-Arg-Gln-Trp-Leu-Ala-Ala-Arg-Ala) (SEQ ID NO. 14), insulin-like growth factor I (57-70) (Ala-Leu-Leu-Glu-Thr-Tyr-Cys-Ala-Thr-Pro-Ala-Lys-Ser-Glu) (SEQ ID NO. 15), insulin-like growth factor I (30-41) (Gly-Tyr-Gly-Ser-Ser-Ser-Arg-Arg-Ala-Pro-Gln-Thr) (SEQ ID NO. 16), insulin-like growth factor I (24-41) (Tyr-Phe-Asn-Lys-Pro-Thr-Gly-Tyr-Gly-Ser-Ser-Ser-Arg-Arg-Ala-Pro-Gln-Thr) (SEQ ID NO. 17), insulin-like growth factor II (33-40) (Ser-Arg-Val-Ser-Arg-Arg-Ser-Arg) (SEQ ID NO. 18), insulin-like growth factor II (33-40) (Tyr-Ser-Arg-Val-Ser-Arg-Arg-Ser-Arg) (SEQ ID NO. 19), insulin-like growth factor II (69-84) (Asp-Val-Ser-Thr-Pro-Pro-Thr-Val-Leu-Pro-Asp-Asn-Phe-Pro-Arg-Tyr) (SEQ ID NO. 20), growth hormone (GH)-releasing peptide-6 (GHRP-6) (His-DTrp-Ala-Trp-DPhe-Lys-NH₂) (SEQ ID NO. 21), beta-Interleukin I (163-171) (Val-Gln-Gly-Glu-Glu-Ser-Asn-Asp-Lys) (SEQ ID NO. 22), beta-Interleukin II (44-56) (Ile-Leu-Asn-Gly-Ile-Asn-Asn-Tyr-Lys-Asn-Pro-Lys-Leu) (SEQ ID NO. 23), Interleukin II (60-70) (Leu-Thr-Phe-Lys-Phe-Tyr-Met-Pro-Lys-Lys-Ala) (SEQ ID NO. 24), exendin-4 (GLP-1 analog) (His-Gly-Glu-Gly-Thr-Phe-Thr-Ser-Asp-Leu-Ser-Lys-Gln-Met-Glu-Glu-Glu-Ala-Val-Arg-Leu-Phe-Ile-Glu-Trp-Leu-Lys-Asn-Gly-Gly-Pro-Ser-Ser-Gly-Ala-Pro-Pro-Pro-Ser-NH₂) (SEQ ID NO. 25), exendin-3 (GLP-1 analog) (His-Ser-Asp-Gly-Thr-Phe-Thr-Ser-Asp-Leu-Ser-Lys-Gln-Met-Glu-Glu-Glu-Ala-Val-Arg-Leu-Phe-Ile-Glu-Trp-Leu-Lys-Asn-Gly-Gly-Pro-Ser-Ser-Gly-Ala-Pro-Pro-Pro-Ser) (SEQ ID NO. 26), epidermal growth factor (20-31) Cys(Acm)-Met-His-Ile-Glu-Ser-Leu-Asp-Ser-Tyr-Thr-Cys(Acm) (SEQ ID NO. 27), bivalirudin (Hirulog) (D-Phe-Pro-Arg-Pro-(Gly)₄-Asn-Gly-Asp-Phe-Glu-Glu-Ile-Pro-Glu-Glu-Tyr-Leu) (SEQ ID NO. 28), hirulog-1 D-Phe-Pro-Arg-Pro-(Gly)₄-Asn-Gly-Asp-Phe-Glu-Glu-Ile-Pro-Glu-Tyr-Leu (SEQ ID NO. 29), C-type natriuretic peptide (1-53) (CNP) (Asp-Leu-Arg-Val-Asp-Thr-Lys-Ser-Arg-Ala-Ala-Trp-Ala-Arg-Leu-Leu-Gln-Glu-His-Pro-Asn-Ala-Arg-Lys-Tyr-Lys-Gly-Ala-Asn-Lys-Lys-Gly-Leu-Ser-Lys-Gly-Cys-Phe-Gly-Leu-Lys-Leu-Asp-Arg-Ile-Gly-Ser-Met-Ser-Gly-Leu-Gly-Cys; Disulfide bridge: Cys37-Cys53) (SEQ ID NO. 30), "Mini ANP" (Met-Cys-His-cyclohexylAla-Gly-Gly-Arg-Met-Asp-Arg-Ile-Ser-Cys-Tyr-Arg, disulfide bridge cys2-cys13) (SEQ ID NO. 31), Melanotan-II (MT-II, alpha-MSH4-10-NH₂, or Ac-Nle4-Asp5-His6-D-Phe7-Arg8-Trp9-Lys10) (SEQ ID NO. 32), thymosin alpha1 (TA1) (Ac-Ser-Asp-Ala-Ala-Val-Asp-Thr-Ser-Ser-Glu-Ile-Thr-Thr-Lys-Asp-Leu-Lys-Glu-Lys-Lys-Glu-Val-Val-

EQ
Conclusion

Glu-Glu-Ala-Glu-Asn) (SEQ ID NO. 33), Cys-Phe-Ile-Gln-Asn-Cys-Pro-Orn-Gly-NH₂, Disulfide bridge: Cys1-Cys6) (SEQ ID NO. 34), octreotide (201-995) (DPhe-Cys-Phe-DTrp-Lys-Thr-Cys-Thr-ol; disulfide bridge: Cys2-Cys7) (SEQ ID NO. 35), calcitonin gene-related peptide (CGRP) (Ala-Cys-Asp-Thr-Ala-Thr-Cys-Val-Thr-His-Arg-Leu-Ala-Gly-Leu-Leu-Ser-Arg-Ser-Gly-Gly-Val-Val-Lys-Asn-Asn-Phe-Val-Pro-Thr-Asn-Val-Gly-Ser-Lys-Ala-Phe-NH₂; Disulfide bridge: Cys2-Cys7) (SEQ ID NO. 36), endomorphin-1 Tyr-Pro-Trp-Phe-NH₂ (SEQ ID NO. 37); endomorphin-2 Tyr-Pro-Phe-Phe-NH₂ (SEQ ID NO. 38), nociceptin (also known as Orphanin FQ, Phe-Gly-Gly-Phe-Thr-Gly-Ala-Arg-Lys-Ser-Ala-Arg-Lys-Leu-Ala-Asn-Gln) (SEQ ID NO. 39), angiotensinogen (1-13) (Asp-Arg-Val-Tyr-Ile-His-Pro-Phe-His-Leu-Val-Ile-His) (SEQ ID NO. 40), adrenomedullin (1-12) (Tyr-Arg-Gln-Ser-Met-Asn-Asn-Phe-Gln-Gly-Leu-Arg) (SEQ ID NO. 41), antiarrhythmic peptide (AAP) (Gly-Pro-Hyp-Gly-Ala-Gly) (SEQ ID NO. 42), Antagonist G (Arg-DTrp-(nMe)Phe-DTrp-Leu-Met-NH₂), indolicidin (Ile-Leu-Pro-Trp-Lys-Trp-Pro-Trp-Trp-Pro-Trp-Arg-Arg-NH₂) (SEQ ID NO. 43), osteocalcin (37-49) (Gly-Phe-Gln-Glu-Ala-Tyr-Arg-Arg-Phe-Tyr-Gly-Pro-Val) (SEQ ID NO. 44), cortistatin 29 (1-13) (Glp)-Glu-Arg-Pro-Pro-Leu-Gln-Gln-Pro-Pro-His-Arg-Asp) (SEQ ID NO. 45), cortistatin 14 Pro-Cys-Lys-Asn-Phe-Phe-Trp-Lys-Thr-Phe-Ser-Ser-Cys-Lys; Disulfide bridge: Cys2-Cys13 (SEQ ID NO. 46), PD-145065 (Ac-D-Bhg-Leu-Asp-Ile-Ile-Trp) (SEQ ID NO. 47), PD-142893 (Ac-D-Dip-Leu-Asp-Ile-Ile-Trp) (SEQ ID NO. 48), fibrinogen binding inhibitor peptide (His-His-Leu-Gly-Gly-Ala-Lys-Gln-Ala-Gly-Asp-Val) (SEQ ID NO. 49), leptin (93-105) (Asn-Val-Ile-Gln-Ile-Ser-Asn-Asp-Leu-Glu-Asn-Leu-Arg) (SEQ ID NO. 50), GR 83074 (Boc-Arg-Ala-DTrp-Phe-DPro-Pro-Nle-NH₂) (SEQ ID NO. 51) Tyr-W-MIF-1 (Tyr-Pro-Trp-Gly-NH₂) (SEQ ID NO. 52), parathyroid hormone related peptide (107-111) (Thr-Arg-Ser-Ala-Trp) (SEQ ID NO. 53), angiotensinogen (1-14) Asp-Arg-Val-Tyr-Ile-His-Pro-Phe-His-Leu-Val-Ile-His-Asn (SEQ ID NO. 54), Leupeptin (Ac-Leu-Leu-Arg-CHO); and further wherein Z comprises at least two identical amino acid units.

Kindly add the following new claims 73-77.